arXiv:1611.09350v1 [math.GR] 26 Nov 2016

A NOTE ABOUT THE $(2,3)$-GENERATION OF $SL_{12}(q)$

TSANKO RAYKOV GENCHEV

Abstract

In this note we provide explicit uniform type $(2,3)$-generators for the special linear group $SL_{12}(q)$ for all q except for $q = 2$ or $q = 4$. Our considerations are easily traceable, self-contained and based only on the known list of maximal subgroups of this group.

Key words: (2,3)-generated group.

2010 Mathematics Subject Classification: 20F05, 20D06.

Now, when the problem concerning the $(2,3)$-generation (especially) of the special linear groups and their projective images is completely solved (see [4] and references there), we give our contribution by discussing the last remaining group $SL_{12}(q)$ in the light of the method used in our works [2], [3] and [5]. It has been proved in [4] the $(2,3)$-generation of the same group over all finite fields in characteristic not equal to 5. The author's approach is different from ours in which we make an essential use of the known list of maximal subgroups of $SL_{12}(q)$.

The group $G = SL_{12}(q)$, where $q = p^m$ and p is a prime, acts naturally on a twelve-dimensional vector space V over the field $F = GF(q)$. We identify V with the column vectors of F^{12}, and let $v_1, \ldots, v_{12}$ be the standard base of the space V. Set $Q = q^{11} - 1$ if $q \neq 3, 7$ and $Q = (q^{11} - 1)/2$ if $q = 3, 7$.

First of all, based only to the orders of the maximal subgroups of G (which exact structure is given in Tables 8.76 and 8.77 in [1]), it can be easily seen that there is only one class of such subgroups having an element of order Q. Namely, this is the class of reducible on the space V subgroups; actually they are the stabilizers in G of one or eleven-dimensional subspaces of V.

Now, let us choose an element ω of order Q in the multiplicative group of the field $GF(q^{11})$ and set

$$f(t) = (t-\omega)(t-\omega^q)(t-\omega^{q^2})(t-\omega^{q^3})(t-\omega^{q^4})(t-\omega^{q^5})(t-\omega^{q^6})(t-\omega^{q^7})(t-\omega^{q^8})(t-\omega^{q^9})(t-\omega^{q^{10}}) = t^{11} - \alpha_1 t^{10} + \alpha_2 t^9 - \alpha_3 t^8 + \alpha_4 t^7 - \alpha_5 t^6 + \alpha_6 t^5 - \alpha_7 t^4 + \alpha_8 t^3 - \alpha_9 t^2 + \alpha_{10} t - \alpha_{11}.$$

Then $f(t) \in F[t]$ and the polynomial $f(t)$ is irreducible over the field F. Note that $\alpha_{11} = \omega^{\frac{q^{11}-1}{q-1}}$ has order $q-1$ if $q \neq 3, 7$, $\alpha_{11} = 1$ if $q = 3$, and $\alpha_{11}^3 = 1 \neq \alpha_{11}$ if $q = 7$.

Set

$$x = \begin{bmatrix} -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_6\alpha_{11}^{-1} & 0 & \alpha_6 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_5\alpha_{11}^{-1} & 0 & \alpha_5 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & \alpha_4\alpha_{11}^{-1} & 0 & \alpha_7 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & \alpha_3\alpha_{11}^{-1} & 0 & \alpha_9 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & \alpha_8\alpha_{11}^{-1} & 0 & \alpha_8 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_7\alpha_{11}^{-1} & 0 & \alpha_4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_1\alpha_{11}^{-1} & -1 & \alpha_{10} \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & \alpha_9\alpha_{11}^{-1} & 0 & \alpha_3 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & \alpha_2\alpha_{11}^{-1} & 0 & \alpha_2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_{11} \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & \alpha_{10}\alpha_{11}^{-1} & 0 & \alpha_1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_{11}^{-1} & 0 & 0 \end{bmatrix},$$

$$y = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}.$$

Then x and y are elements of G of orders 2 and 3, respectively. Denote

$$z = xy = \begin{bmatrix} 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_6 & \alpha_6\alpha_{11}^{-1} \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_5 & \alpha_5\alpha_{11}^{-1} \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & \alpha_7 & \alpha_4\alpha_{11}^{-1} \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & \alpha_9 & \alpha_3\alpha_{11}^{-1} \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_8 & \alpha_8\alpha_{11}^{-1} \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_4 & \alpha_7\alpha_{11}^{-1} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & \alpha_{10} & \alpha_1\alpha_{11}^{-1} \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & \alpha_3 & \alpha_9\alpha_{11}^{-1} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & \alpha_2 & \alpha_2\alpha_{11}^{-1} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_{11} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & \alpha_1 & \alpha_{10}\alpha_{11}^{-1} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \alpha_{11}^{-1} \end{bmatrix}.$$

The characteristic polynomial of z is $f_z(t) = (t - \alpha_{11}^{-1})f(t)$ and the characteristic roots α_{11}^{-1}, ω, ω^q, ω^{q^2}, ω^{q^3}, ω^{q^4}, ω^{q^5}, ω^{q^6}, ω^{q^7}, ω^{q^8}, ω^{q^9}, and $\omega^{q^{10}}$ of z are pairwise distinct. Then, in $GL_{12}(q^{11})$, z is conjugate to the matrix diag (α_{11}^{-1}, ω, ω^q, ω^{q^2}, ω^{q^3}, ω^{q^4}, ω^{q^5}, ω^{q^6}, ω^{q^7}, ω^{q^8}, ω^{q^9}, $\omega^{q^{10}}$) and hence z is an element of $SL_{12}(q)$ of order Q.

Let $H = \langle x, y\rangle$, $H \leq G$. We prove that H acts irreducibly on the space V. Indeed, assume that W is an H-invariant subspace of V and $k = \dim W$, $1 \leq k \leq 11$.

Let first $k = 1$ and $0 \neq w \in W$. Then $y(w) = \lambda w$ where $\lambda \in F$ and $\lambda^3 = 1$. This yields

$$w = \mu_1(v_1+\lambda^2 v_2+\lambda v_3)+\mu_2(v_4+\lambda^2 v_5+\lambda v_6)+\mu_3(v_7+\lambda^2 v_8+\lambda v_9)+\mu_4(v_{10}+\lambda^2 v_{11}+\lambda v_{12}),$$

where μ_1, μ_2, μ_3, and μ_4 are elements of the field F.
Now, we involve the action of x onto w: $x(w) = \nu w$ where $\nu = \pm 1$. This yields consecutively $\mu_4 \neq 0$, $\alpha_{11} = \lambda^2\nu$, and

$$\mu_3 = \lambda(\alpha_1 + \nu\alpha_{10} - \lambda\nu)\mu_4, \tag{1}$$

$$\nu\mu_1 + \mu_2 = (\nu\alpha_4 + \alpha_7)\mu_4, \tag{2}$$

$$\nu\mu_2 + \lambda^2\mu_3 = \lambda(\nu\alpha_3 + \alpha_9)\mu_4, \tag{3}$$

$$(\nu + 1)(\mu_1 - \lambda\alpha_6\mu_4) = 0, \tag{4}$$

$$(\nu + 1)(\mu_1 - \lambda^2\alpha_5\mu_4) = 0, \tag{5}$$

$$(\nu + 1)(\mu_2 - \lambda^2\alpha_8\mu_4) = 0, \tag{6}$$

$$(\nu + 1)(\mu_3 - \alpha_2\mu_4) = 0, \tag{7}$$

In particular, we have $\alpha_{11}^3 = \nu$ and $\alpha_{11}^6 = 1$. This is impossible if $q = 5$ or $q > 7$ since then α_{11} has order $q - 1$. According to our assumption ($q \neq 2, 4$) only two possibilities left: $q = 3$ (and $\alpha_{11} = 1$) or $q = 7$ (and $\alpha_{11}^3 = 1 \neq \alpha_{11}$). So $\nu = 1$, $\alpha_{11} = \lambda^2$ and (1), (2), (3), (4), (5), (6) and (7) produce $\alpha_1 = \lambda^2\alpha_2 - \alpha_{10} + \lambda$, $\alpha_6 = \lambda\alpha_5$, $\alpha_7 = \lambda^2\alpha_5 + \lambda^2\alpha_8 - \alpha_4$ and $\alpha_9 = \lambda\alpha_2 + \lambda\alpha_8 - \alpha_3$. Now $f(-1) = -(1 + \lambda + \lambda^2)(1 + \alpha_2 + \alpha_5 + \alpha_8) = 0$ both for $q = 3$ and $q = 7$, an impossibility as $f(t)$ is irreducible over the field F.

Now let $2 \leq k \leq 11$. Then the characteristic polynomial of $z|_W$ has degree k and has to divide $f_z(t) = (t - \alpha_{11}^{-1})f(t)$.The irreducibility of $f(t)$ over F leads immediately to the conclusion that this polynomial is $f(t)$ and $k = 11$. Now the subspace U of V which is generated by the vectors v_1, v_2, v_3, . . . , v_{11} is $\langle z\rangle$-invariant. If $W \neq U$ then $U \cap W$ is $\langle z\rangle$-invariant and dim $(U \cap W) = 10$. This means that the characteristic polynomial of $z|_{U\cap W}$ has degree 10 and must divide $f_z(t) = (t - \alpha_{11}^{-1})f(t)$ which is impossible. Thus $W = U$ but obviously U is not $\langle y\rangle$-invariant, a contradiction.

Note that the above considerations fail if $q = 2$ or 4.

Now, as $H = \langle x, y\rangle$ acts irreducible on the space V and it has an element of order Q, we conclude that H can not be contained in any maximal subgroup of $G(= SL_{12}(q))$. Thus $H = G$ and $G = \langle x, y\rangle$ is a $(2, 3)$-generated group. □

References

[1] J. N. BRAY, D.F. HOLT, C. M. RONEY-DOUGAL. The Maximal Subgroups of the Low - Dimensional Finite Classical Groups. London Math. Soc. Lecture Note Series **407**, Cambridge University Press (2013).

[2] TS. GENCHEV, E. GENCHEVA. $(2,3)$-generation of the special linear groups of dimension 8. *Proceedings of the Forty Fourth Spring Conference of the Union of Bulgarian Mathematicians, SOK "Kamchia"*, April 2-6 (2015), 167-173.

[3] E. GENCHEVA, TS. GENCHEV and K. TABAKOV. $(2,3)$-generation of the special linear groups of dimensions 9, 10 and 11. *arXiv* **1412.8631v5** (2016).

[4] M. A. PELLEGRINI. The $(2,3)$-generation of the special linear groups over finite fields. *arXiv* **1605.04276v1** (2016).

[5] K. TABAKOV, E. GENCHEVA and TS. GENCHEV. $(2,3)$-generation of the special linear groups of dimension 11. *Proceedings of the Forty Fifth Spring Conference of the Union of Bulgarian Mathematicians, Pleven*, April 6-10 (2016), 146-151.

Ts. Genchev
e-mail: genchev57@yahoo.com
Department of Mathematics
Technical University
Varna, Bulgaria